

Nota Científica

Description of *Isomeria incomparabilis* nov. spec. (Gastropoda: Labyrinthidae) from Northern Ecuador

Descripción de *Isomeria incomparabilis* nov. spec. (Gastropoda: Labyrinthidae) para el norte de Ecuador

Marijn T. Roosen^{1,2,3,*}   , Carles Dorado⁴  

¹ Instituto Nacional de Biodiversidad (INABIO), Pje. Rumipamba N. 341 y Av. de los Shyris (Parque La Carolina), Quito, Ecuador

² Naturalis Biodiversity Centre, P.O. Box 9517, 2300 RA Leiden, The Netherlands

³ Natural History Museum of Rotterdam, Westzeedijk 345, 3015 AA Rotterdam, The Netherlands

⁴ Associació Catalana de Malacología, Museu Blau, Plaça Leonardo da Vinci 4-5, 08019 Barcelona, Spain

DOI: <https://doi.org/10.26807/remcb.v46i1.1051> | ***Autor de correspondencia:** roosen@hetnatuurhistorisch.nl
Fecha de envío: 04/05/2025 | **Fecha de aceptación:** 28/10/2025 | **Fecha de publicación:** 21/11/2025

Abstract.- A new species, *Isomeria incomparabilis* nov. spec., is described and tentatively assigned to *Isomeria* Albers, 1850. No similar species are known.

Keywords: *Isomeria* Albers, 1850, *Labyrinthus* Beck, 1837, Imbabura, new species, Andes

ZooBank ID: urn:lsid:zoobank.org:pub:61CBEA4D-D580-4E7F-B861-FC3E878065E6

Resumen.- En el presente artículo se describe *Isomeria incomparabilis* nov. spec. la cual se asigna tentativamente a *Isomeria* Albers, 1850. No se conocen especies similares.

Palabras Clave: *Isomeria* Albers, 1850, *Labyrinthus* Beck, 1837, Imbabura, especie nueva, Andes

ZooBank ID: urn:lsid:zoobank.org:pub:61CBEA4D-D580-4E7F-B861-FC3E878065E6

Introduction

Sometimes terrestrial gastropod species that do not resemble any other taxa from the country are still found in Ecuador. Examples of this are the recently described species of *Adelopoma* Doering, 1885 and *Xenodiscula* Pilsbry, 1919 (Greke et al. 2023; Roosen et al. 2023). Usually these completely unknown species are small, but sometimes larger species are discovered.

In this paper we describe a new species of Labyrinthidae (Borrero et al. 2017), which we assign to *Isomeria* Albers, 1850 because of its apertural dentition. Its small and strongly carinated shell with colour bands separate it from other species of *Isomeria*. The type material of *I. incomparabilis* nov. spec. is insufficient to investigate its affinities.



Como citar este artículo: Roosen MT, Dorado C. 2025. Description of *Isomeria incomparabilis* nov. spec. (Gastropoda: Labyrinthidae) from Northern Ecuador. Revista Ecuatoriana de Medicina y Ciencias Biológicas 46(2): 115-118. doi: 10.26807/remcb.v46i2.1051



Material and Methods

The holotype of *I. incomparabilis* nov. spec. is present in the collection of Pontificia Universidad Católica del Ecuador (Quito-Ecuador) (PUCE, QCAZI). Locality data was recorded from the original label. The shell was imaged with SONY DSC-HX60V camera by the second author (CD). To confirm *I. incomparabilis* nov. spec. was new to science; it was compared to all Labyrinthidae in Breure et al. (2022) and Solem (1966). Similar species will be compared to it based on the images in these publications. Whorls were counted to the nearest 1/4th whorl using the method of Gittenberger et al. (1998).

Results

Class Gastropoda Cuvier, 1795

Order Stylommatophora A. Schmidt, 1855

Family Labyrinthidae Borrero, Sei, D. G. Robinson & Rosenberg, 2017

Genus *Isomeria* Albers, 1850

Isomeria Albers, 1850: 126. Type species (by monotypy): *Helix oreas* F.C.L. Koch, 1844

Isomeria incomparabilis nov. spec.

Zoobank ID: urn:lsid:zoobank.org:act:23E4DCD3-0A59-4A05-B8D2-FC9ABC2DB31F

Type material.- QCAZI 278700 (holotype, one shell, dry), Leg. M. Navarrete, 02-11-2012.

Type locality. - Ecuador, Imbabura Province, Lita. (00°52'21,77"N, 78°28'00,60"W), 780 meters above sea level.

Measurements.- 28.12 mm (W), 12.83 mm (H).

Description.- Shell medium large, but small for the genus; strongly carinate; strongly depressed conical in shape. No clear differences visible between the protoconch and the teleoconch. The sculpture consists of indistinct growth lines combined with a granulated surface. The granulations are limited to the section between the carina and the umbilical wall. In addition, some sections above the carina and on the umbilical wall show minute spiral grooves. Aperture trapezoid, slightly deflected, with a thickened peristome. Two small palatal teeth are present, one almost in the carina. Umbilicus wide, about 17% of total width. Colour of the shell is light, hazelnut brown, with three dark, chestnut brown bands located below the suture, around the carina and in the umbilicus. The carina itself is white (Figure 1).

Geographic range.- Ecuador. Only known from the type locality in the Imbabura Province.

Habitat.- The specimen was found fixed on a trunk, implying a ground dwelling lifestyle.

Comparisons.- No similar species are known from Ecuador. It is reminiscent of *Isomeria minuta* Solem, 1966, but this species has a different dentition (one basal tooth and one palatal tooth), smaller size and does not have any distinct colour patterns. Another species with which it shares some characters is *Labyrinthus dunkeri* (L. Pfeiffer, 1852), which has a similar colour pattern, carinated shell and granulated surface. However, *L. dunkeri* differs from the new species by its smaller size, 18.6 - 24.9 mm wide according to Solem (1966), and more complex apertural dentition which include one parietal lamella and two basal teeth.

Etymology.- The specific epithet *incomparabilis* refers to the difficulty in finding a similar species to compare it with.

Remarks.- Based on the limited apertural dentition, we describe *I. incomparabilis* nov. spec. as a species of *Isomeria*, even though other shell characters most closely resemble a species currently included in *Labyrinthus* (*L. dunkeri*). More, preferably living, specimens need to be found and studied to confirm this placement.

Isomeria incomparabilis nov. spec. resembles *Labyrinthus* Beck (1837) due to its less inflated, carinated whorls and low spire, but *Labyrinthus* species have more numerous and/or larger teeth and lamella on the palatal and parietal wall (Borrero et. al. 2017). Moreover, in Ecuador *Labyrinthus* only occurs in the Amazon rainforest and on the eastern slopes of the Andes (Breure et al. 2022; Solem 1966). It is possible that apertural dentition should not always be the main character to distinguish between these genera, but we do not have the information needed to study this. Sequencing and comparing the DNA of *I. incomparabilis* nov. spec. and *L. dunkeri* could help answer this question.

In the description of *Isomeria minuta*, Solem (1966) also mentions that this species shows characters referable to both *Isomeria* and *Labyrinthus* and highlights that the transition between both shell shapes may have happened several times during their evolution.

For all papers we published the past years we visited several main museums in Ecuador and several countries in Europe (The Netherlands, Belgium, Spain, etc.). We did not find this species in any other collections.



Figure 1. *Isomeria incomparabilis* nov. spec. Holotype (QCAZI 278700), dry shell, from Lita, Imbabura Province, northwest Ecuador. Scale 10 mm.

Discussion

This paper describes yet another new species of terrestrial gastropod from Ecuador. This species does not teach us a lot at the moment. Of course, seen in context it shows that even larger gastropod species have yet to be discovered in Ecuador, but as it is based on only one empty shell it does not provide additional insights on the ecological relationships of gastropod communities in the rainforests of Ecuador. In any case, its discovery is a significant addition to the country's malacofauna and deserves to be published. That is also the main purpose of this paper: showing that truly aberrant, large species can still be discovered in areas that have received little attention in the past.

Hopefully the description of *I. incomparabilis* nov. spec. will lead to more research into this and other poorly known species. In addition, this description hopefully increases interest in collecting new material in less studied Ecuadorian provinces, such as Carchi and Esmeraldas, that are very close to the type locality.

Acknowledgements

We would like to thank Fernanda Salazar (PUCE-QCAZI) and her colleagues for providing us access to their invertebrate collections, which have been of great value for the development of this paper.

Declaration of conflict of interest

The authors declare no conflict of interest.

Authors' contribution

MR and CD recognised the species as new, compared it to its congeners and wrote the manuscript.

References

Breure ASH, Roosen MT, Ablett JD. 2022. Land and freshwater molluscs of mainland Ecuador: an illustrated checklist. *Iberus*. 40:1-290.

Greke K, Kagainis U, Telnov D. 2023. First record of the family Diplommatinidae Gray, 1847 (Gastropoda: Architaenioglossa) from Ecuador with description of a new *Adelopoma* Doering, 1885. *Zootaxa*. 5339(3):273-284. doi:10.11646/zootaxa.5339.3.4.

Gittenberger E, Janssen AW, Kuijper WJ, Kuiper JGJ, Meijer T, Van der Velde G, De Vries JN. 1998. De Nederlandse zoetwatermollusken: recente en fossiele weekdieren uit zoet en brak water. Leiden (NL): Naturalis Biodiversity Center.

Roosen MT, Weijsenfeld J, Dorado C. 2023. Notes on the genus *Xenodiscula* Pilsbry, 1919 (Gastropoda: Scolodontidae), with the description of a new species from NW Ecuador. *Folia Malacol.* 31(2):100-106. doi:10.12657/folmal.031.014.

Sei M, Robinson DG, Geneva AJ, Rosenberg G. 2017. Doubled helix: Sagdoidea is the overlooked sister group of Helicoidea (Mollusca: Gastropoda: Pulmonata). *Biol J Linn Soc.* 122(4):697-728. doi:10.1093/biolinnean/blx082.

Solem A. 1966. The neotropical land snail genera *Labyrinthus* and *Isomeria* (Pulmonata, Camaenidae). *Fieldiana Zool.* 50:1-226. Available from: <https://www.biodiversitylibrary.org/page/2780997#page/15/mode/1up>