

Artículo científico

Diversity of fungal communities inhabiting symptomatic leaves of Cape gooseberry (*Physalis peruviana*) in the Ecuadorian Andes

Diversidad de comunidades fúngicas que habitan en tejido foliar sintomático de la uvilla *Physalis peruviana* en los Andes Ecuatorianos

José Luis Díaz¹, Wagner Chaves-Acuña^{1*} & Jeniffer Yáñez¹

¹ Escuela de Ciencias Biológicas, Pontificia Universidad Católica del Ecuador, Avenida 12 de Octubre 1076 y Roca, Apartado 17-01-2184, Quito, Ecuador

*wchaves512@gmail.com

<https://doi.org/10.26807/remcb.v40i2.762>

Recibido 30-05-2019 ; Aceptado 24-09-2019

ABSTRACT.- We used microbial culture specific methods to assess the occurrence and diversity of fungal communities inhabiting symptomatic foliar tissue of Cape gooseberry (*Physalis peruviana*) at four different field plots in the northern Ecuadorian Andes. We categorized the isolated fungi based on their morphological characteristics and subsequently identified them through rDNA sequencing of the ITS region. We recovered 75 ascomycetous isolates which represented 13 genera, revealing a total of 20 taxa. We found most isolates recovered from necrotic and straw-colored tissues to belong to the genus *Alternaria*. Other fungal taxa frequently found were *Epicoccum*, *Diaporthe*, and *Xylaria*. To our knowledge, this is the first study to report the mycobiota associated to *P. peruviana* plants showing disease symptoms, including 2 new genera of fungi for the first time in Ecuador. Our results are expected to provide useful information for future assessments of biological control on tropical commercial.

KEYWORDS: Ascomycota, rDNA, ITS, plant disease, uvilla

RESUMEN.- Utilizamos métodos cultivo-dependientes para evaluar la ocurrencia y diversidad de comunidades fúngicas que habitan en tejido foliar sintomático de uvilla (*Physalis peruviana*) en cuatro localidades del norte de los Andes ecuatorianos. Categorizamos los aislados basándonos en las características morfológicas de los cultivos obtenidos *in vitro*, los cuales identificamos al secuenciar la región de ADNr ITS. En total, recuperamos un total de 75 hongos ascomycetos que comprendían 13 géneros y 20 taxa. Encontramos que la mayoría de hongos recuperados de tejidos necróticos o cloróticos pertenecían al género *Alternaria*. Otros taxa que encontramos con frecuencia fueron *Epicoccum*, *Diaporthe* y *Xylaria*. A nuestro conocimiento, este es el primer estudio en reportar la micobiota asociada a plantas de *P. peruviana* que presenten hojas con síntomas de enfermedad, incluyendo 2 géneros de hongos nuevos para el Ecuador. Esperamos que nuestros resultados provean información útil para futuros estudios sobre control biológico aplicado a cultivos comerciales en los trópicos.

PALABRAS CLAVE: Ascomycota, ADNr, ITS, patología vegetal, uvilla

INTRODUCTION

Cape gooseberry (*Physalis peruviana* L.) is a biannual shrub native from the Andean highlands of northern South America that can grow up to 1 m

high (Legge 1974; Fischer 2000). In recent years, Cape gooseberries have gained increasing attention in the markets of South and Central America, due to the plant's high tolerance to a broad variety of environments (Ramadan and Moersel 2003; Puente

et al. 2011). Given the variety of antioxidant types and high sugar content (Novoa et al. 2006), this crop is of great economic importance for industrial purposes worldwide (Morton 1987; Rehm and Espig 1991; McCain 1993; Ramadan and Moersel 2003; Mazorra 2006), as it has also been widely used for its medicinal properties to treat cancer and diabetes (see review in Puente et al. 2011).

In Ecuador, *P. peruviana*, locally known as *uvilla*, is grown in the northern region between 2 000 – 3 000 m.a.s.l., where moderately cold temperatures prevail (Puente et al. 2011; Fischer et al. 2014). The production of *P. peruviana* has increased considerably in recent years across Ecuador due to its growing export demands to European countries (Muñoz 2003; Altamirano 2010; Fischer et al. 2014). However, a wide array of selective pressures including biotic factors (herbivory, parasites) and abiotic stress (low nutrient availability, drought) may result in major production losses by inducing yellowing or necrosis in photosynthetic tissues (Douanla-Meli et al. 2013). As a result, several toxic pesticides (e.g. herbicides and fungicides) have been used in this region as agricultural practices to prevent plant diseases (Rodríguez-Amézquita et al. 2010). Yet, even though the main biological constraint to Cape gooseberry are fungi of the genera *Cladosporium*, *Phoma*, *Alternaria*, *Botrytis* and *Colletotrichum* (Angulo 2005; Fischer and Miranda 2012; De La-Rotta 2014), little is known on the mycobiota inhabiting symptomatic leaves of *P. peruviana*, which could provide useful information and assist crop management.

Studies based solely on morphology have failed to be conclusive at taxonomic species level for fungi associated to *P. peruviana* (Crozier et al. 2006), while fungal isolates often fail to form fruiting structures in culture (Gazis and Chaverri 2010). Thus, considering that leaves, as ecological niches, might influence the diversity and composition of fungal symbionts (Kriel et al. 2000), the present study is aimed at understanding the community structure and diversity of fungi isolated from symptomatic leaves of Cape gooseberry in the Andean highlands of northern Ecuador.

MATERIALS AND METHODS

Sample collection.- We collected unhealthy leaf samples showing necrosis symptoms or straw-colored tissues in plantations of *Physalis peruviana* during June 2015 at four different sites of the Pichincha Province in the northern Ecuadorian

Andes: La Merced (00°17.635' S, 078°24.158' W, 2 611 m.a.s.l.), Virgen de Lourdes (00°17.311' S, 078°24.384' W, 2 628 m.a.s.l.), Tumbaco (00°13.539' S, 078°23.979' W, 2 450 m.a.s.l.), and Yaruquí (00°11.596' S, 078°19.710' W, 2 579 m.a.s.l.). We randomly selected 14 mature *P. peruviana* (2 to 7 plants per site > 0.5 m in height) and then transported 75 leaf samples (5 x 5 mm) to the laboratory in sterile plastic bags. We stored the leaves for 2 d at -10 °C before the isolation of fungi.

Isolation of fungi.- To isolate fungi, we followed a modified protocol from Crous et al. (2009). In order to induce sporulation, we surface-disinfected the samples through immersion in 70 % ethanol for 1 min and 3 % sodium hypochlorite for 5 min. We rinsed the samples with sterile distilled water three times before transferring them to Petri dishes (90 mm) that contained a sterile Potato Dextrose Agar (PDA, Difco Laboratories, Detroit, MI) supplemented with streptomycin to suppress bacterial proliferation. We incubated the plates at 25 °C with a 12-h photoperiod and checked them regularly for fungal growth for 7 to 15 days. After fungal growth was visible, we subcultured different colonies on new Petri dishes with the same medium for further purification and identification (Crozier et al. 2006). We categorized and divided filamentous fungi isolates based on the morphology of the colonies.

DNA extraction.- We collected about 10 mg of fresh fungal tissue using a sterile scalpel and extracted genomic DNA using the Wizard Genomic DNA Purification Kit (Promega, Madison, WI) in accordance with the manufacturer's instructions. To each tissue sample, we added 600 µl of nuclear lysis solution and vortexed the resulting solution for 15 min at 65 °C, followed by 15 min of water immersion at 37 °C. Then, we added 3 µl of RNAase solution and incubated the resulting solution at room temperature for 5 min, before we mixed it with a protein precipitation solution. We vortexed the samples for 15 s and centrifuged the protein precipitation solution at 13 000–16 000 g for 3 min. We transferred the supernatant liquid into a 1.5 mL fresh tube mixed with 600 µl ethanol at room temperature and centrifuged the columns at 13 000–16 000 g for 1 min, before we discarded the flow-through. Finally, we washed the samples as per manufacturer's protocol and we quantified the purified DNA concentration with a Qubit® 20 Quantitation Starter Kit (Invitrogen, USA) following instructions provided by the manufacturer and stored the tubes at -20 °C.

PCR amplification.- We used PCR to amplify primer pairs of the Internal Transcribed Spacers (ITS), ITS4 (5'-TCCTCCGCTTATTGATATGC-3') and ITS5 (5'-GGAAGTAAAAGTCGTAACAAGG-3'), which are rDNA genes commonly used for fungal barcoding (White et al. 1990). We performed PCR using GoTaq Green Master mix (Promega, Madison, WI) according to manufacturer's instructions. We carried out reactions in 25 µl volume samples as follows: first initial denaturation at 94 °C for 5 min, followed by 30 cycles at 94 °C for 30 s, 55 °C for 30 s, 72 °C for 30 s, and a final elongation at 72 °C for 10 min in a mixture of 22 µl GoTaq® Green Master Mix, 10 µM of primers and 2 µl of template. We analyzed PCR products by electrophoresis on a 2,5 % (w/v) agarose gel containing Sybr Safe (Invitrogen, Mulgrave, Australia) and visualized them under UV light. We purified the samples using the Wizard SV Gel and PCR Clean-up System (Promega). Reaction mixtures were sequenced by a private biotechnology company (Macrogen, Seoul, South Korea).

Fungal identification.- To identify the fungal isolates based on ITS, we compared nucleotide sequences against GenBank's database using the basic local alignment search tool (BLASTn) with the National Center for Biotechnology Information (NCBI, Bethesda, MD) database. Whenever possible, we defined fungi species considered in this study using an ITS similarity threshold of 97 % to sequences deposited in GenBank (Nilsson et al. 2008). If the ITS sequence was not discriminant at a species level, we assigned the isolates to the closest taxonomic unit. We deposited all gene sequences retrieved in this work in the GenBank database under the accession numbers: MF434350.1–MF435178.1. Fungal classification follows MycoBank (<http://www.mycobank.org>) database.

Statistical analysis.- We measured fungal occurrence considering both, the colonization frequency (CF) and the relative frequency of isolation (RF). We calculated CF as the total number of segments colonized by a given fungus divided by the total number of segments in that sample, expressed as percentage. We determined RF as the total number of segments colonized by a given taxon divided by the total number of taxa. We calculated fungal species dominance, diversity, and richness using Simpson's, Shannon's and Margalef's indices, respectively. We included singlets (occurrence of ≤ 2 isolates) for all analyses as they are likely to be keystone species of symptomatic foliar tissues (Gazis and Chaverri

2010). We calculated all diversity indices with PAST, version 1.9 (Paleontological Statistics) software (Hammer et al. 2001), in accordance to the following formulas:

1. Simpson's diversity ($1-D$) = $1 - \sum (p_i)^2$
2. Shannon-Wiener index (H') = $-\sum p_i (\log_2 p_i)$
3. Margalef's (D^{mg}) = $(S-1) / \ln N$

where, p_i is the proportion of colonization frequency of a given taxon, S is the total number of taxa isolated, and N is the number of species recorded.

RESULTS

Identification by ITS rDNA sequencing showed that all fungal isolates recovered in this study were Ascomycota, placed in seven orders: Pleosporales, Xylariales, Capnodiales, Glomerellales, Dothideales, Diaporthales and Hypocreales. Most isolates belonged to the Pleosporales (40 %) and Xylariales (20 %) orders. BLAST analysis revealed 13 different genera associated to symptomatic tissues of *P. peruviana*, while we were not able to identify the genus of two isolates and limited their categorization to the order level (Table 1). From a total of 20 identified taxa, the most frequently recovered fungal genera were *Alternaria*, *Epicoccum* and *Diaporthe* (Table 2), while 20 % of the isolates occurred as doubletons (occurrence of 2 isolates) and 50 % occurred only once. Fungal communities showed a low overall isolation rate of 5 ± 7.50 %, while diversity indices displayed high values for species dominance ($1-D = 0,85$), diversity ($H' = 3,33$), and richness ($D^{mg} = 4,40$).

DISCUSSION

To date, almost all species in the plant kingdom are known to host bacteria or fungi within the intra and inter-cellular spaces of their vegetative tissues (Petrini 1996; Arnold and Lutzoni 2007; Mondal et al. 2007; Hyde and Soyong 2008, Ghimire et al. 2011, Rocha et al. 2011, Douanla-Meli et al. 2013). Despite the fact that there is no apparent harm caused by these microorganisms on the plant's health, many fungal species have been reported to enter the plant through stomata or wounds (Andersen and Walker 1985; Schulz and Boyle 2006; Sieber 2007), while some species may act as latent pathogens during the host's senescence (Photita et al. 2004; Promputtha et al. 2007; Rodríguez and Redman 2008; Prihastuti et al. 2009).

Tabla 1. ITS-based identification of fungal isolates recovered from symptomatic leaves of *Physalis peruviana* in the Ecuadorian Andes.

Proposed fungal taxon	^a GenBank Accession No.	QC (%)	Id (%)	Sequence length (bp)	^b Top BLAST search results (GenBank accession No.)	Reference
<i>Alternaria</i> sp.	MF435050.1	100	100	568	KM215624.1	Raja et al. 2015
<i>Bipolaris cynodontis</i>	MF435062.1	100	100	541	KJ909767.1	Manamgoda et al. 2014
<i>Cercospora dubia</i>	MF435168.1	99	100	537	KX287277.1	Videira et al. 2016
<i>Colletotrichum boninense</i>	MF435150.1	100	98.99	592	JX258799.1	Weir et al. 2012
<i>Colletotrichum gloeosporioides</i>	MF435164.1	100	100	571	KM257026.1	Waculicz-Andrade et al. 2017
<i>Diaporthe helianthi</i>	MF435054.1	98	97.73	579	AJ312356.1	Rekab et al. 2004
<i>Diaporthe</i> sp.	MF435071.1	100	99.65	572	KC339218.1	Panno et al. 2013
<i>Didymella glomerata</i>	MF435166.1	100	99.81	538	AY183371.1	Catal 2002
<i>Dothideales</i> sp.	MF435051.1	99	99.64	553	HQ607988.1	Rodrigues et al. 2011
<i>Epicoccum nigrum</i>	MF435053.1	100	100	544	KX869965.1	Kernaghan et al. 2017
<i>Fusarium equiseti</i>	MF435058.1	100	100	547	KR094440.1	Kaur et al. 2016
<i>Fusarium venenatum</i>	MF435064.1	100	100	523	KP295496.1	Stefańczyk et al. 2016
<i>Nigrospora oryzae</i>	MF435092.1	100	100	550	EU272503.1	Miles et al. 2012
<i>Periconia byssoides</i>	MF435088.1	97	99.82	577	MK370654.1	Herrmann et al. 2019
<i>Phoma</i> sp.	MF435059.1	100	99.26	538	JN207257.1	Loro et al. 2012
<i>Pleosporales</i> sp.	MF435097.1	100	99.62	533	HQ631051.1	Shrestha et al. 2011
<i>Stagonosporopsis cucurbitacearum</i>	MF435112.1	100	100	552	GU045304.1	Ling et al. 2010
<i>Xylaria multiplex</i>	MF435085.1	99	99.66	588	KP133436.1	Thomas et al. 2016
<i>Xylaria</i> sp.	MF435158.1	100	98.45	576	FJ799949.1	Van Bael et al. 2009
<i>Xylaria venosula</i>	MF435102.1	97	99.65	587	EF026149.1	Hsieh et al. 2010

^a All sequences were deposited in the NCBI GenBank (Accession No. column). QC: query cover, Id: identity. ^b

Sequences available in GenBank used for comparisons with sequences obtained in this study via BLAST software.

Tabla 2. Total number of isolates, isolation rate (%) and relative frequency of ascomycetous fungi recovered from symptomatic foliar tissues of *Physalis peruviana* in the Ecuadorian Andes.

Taxa	No. Isolates	Isolation rate (%)	Relative frequency
Capnodiales			
<i>Cercospora malayensis</i>	1	0.775	0.0244
Diaporthales			
<i>Diaporthe helianthi</i>	3	2.326	0.0732
<i>Diaporthe</i> sp.	11	8.527	0.2683
Dothideales			
Dothideales sp.	1	0.775	0.0244
Glomerellales			
<i>Colletotrichum boninense</i>	1	0.775	0.0244
<i>Colletotrichum gloeosporioides</i>	1	0.775	0.0244
Hypocreales			
<i>Fusarium equiseti</i>	2	1.550	0.0488
<i>Fusarium venenatum</i>	1	0.775	0.0244
Pleosporales			
<i>Alternaria</i> sp.	22	17.054	0.5366
<i>Bipolaris cynodontis</i>	2	1.550	0.0488
<i>Didymella glomerata</i>	2	1.550	0.0488
<i>Epicoccum nigrum</i>	15	11.628	0.3659
<i>Periconia byssoides</i>	1	0.775	0.0244
<i>Phoma</i> sp.	2	1.550	0.0488
Pleosporales sp.	1	0.775	0.0244
<i>Stagonosporopsis cucurbitacearum</i>	1	0.775	0.0244
Xylariales			
<i>Nigrospora oryzae</i>	3	2.326	0.7320
<i>Xylaria multiplex</i>	3	2.326	0.0732
<i>Xylaria</i> sp.	1	0.775	0.0244
<i>Xylaria venosula</i>	1	0.775	0.0244

In the present study, we used ITS sequencing to identify fungal communities inhabiting symptomatic leaves of *P. peruviana*. Our results are consistent with previous studies that have also shown a dominance by Ascomycota members within fungal assemblages of tropical plants (Arnold and Lutzoni 2007; Gonzaga et al. 2015). Previous studies on fungal assemblages of commercial crops have reported *Alternaria*, *Colletotrichum*, *Epicoccum*, *Fusarium*, *Nigrospora*, *Phoma*, and *Xylaria* taxa inhabiting foliar tissues (Crous et al. 1995; Gazis and Chaverri 2010; Parsa et al. 2016). However, most studies have focused on isolating fungal endophytes inhabiting asymptomatic aerial organs, while very few of them have investigated the fungal species from symptomatic tissues (Maher et al. 2012; Bruez et al. 2014; Dávila et al. 2018). This work constitutes the first report

of mycobiota on Cape gooseberry, from which *Alternaria*, *Bipolaris*, *Cercospora*, *Colletotrichum*, *Didymella*, *Epicoccum*, *Fusarium*, *Nigrospora*, *Phoma* and *Xylaria* taxa have been previously reported in Ecuador (Evans and Reeder 2000; Pacin et al. 2003; Ramírez et al. 2006; Thomas et al. 2008; Cornejo-Espinoza 2014; Moya-Maldonado 2016; Dávila et al. 2018), whereas *Periconia* and *Stagonosporopsis* represent new records for the mycoflora of continental Ecuador and are reported for the first time in *P. peruviana* within its center of origin.

The overall colonization of fungal species screened in the present study is low when compared to that reported in asymptomatic tissues of other plant species (Arnold and Lutzoni 2007; Sun et al. 2008). However, the diversity and richness of species is

similar to other studies of fungal communities inhabiting symptomatic tissues in tropical crops (Dávila et al. 2018). Furthermore, the most dominant genera isolated in the current study were *Alternaria* and *Epicoccum*, which coincides with other fungal communities from tropical angiosperms (Parsa et al. 2016). Within the fungal isolates obtained here, we detected common Cape gooseberry pathogens belonging to the genera *Alternaria*, *Fusarium*, *Colletotrichum*, *Phoma* and *Cercospora* (Rao and Subramonian 1976; Zapata et al. 2002; Angulo 2005; Fischer and Miranda 2012; De La-Rotta 2014) although only the *Alternaria* genus was isolated considerably in this study. As we did not conduct Koch's postulates for the isolates obtained in this study, our results do not imply that the isolated taxa are pathogenic.

The great diversity of fungal species reported in this study is expected to provide novel information for the crop management and biological control on the Ecuadorian Cape gooseberry, when dealing with necrosis symptoms in leaves. The current inventory of fungal species in *P. peruviana* suggests potential sources of culturable secondary metabolites isolated from decaying foliar tissues (Paparú et al. 2008), which may help to establish healthier agricultural practices, while it also contributes to the ecological understanding of specificity patterns shown by fungal communities. Further research of fungal interactions in Cape gooseberry is needed to comprehend the role that the most abundant species isolated in this study play within the fungal assemblages of symptomatic leaf tissues.

ACKNOWLEDGMENTS

This investigation was funded by the Pontificia Universidad Católica del Ecuador. We thank Raúl Jaramillo for his comments and suggestions concerning this research. The authors have no conflict of interest to declare.

REFERENCES

- Altamirano M. 2010. Estudio de la cadena productiva de uvilla (*Physalis peruviana* L.) en la Sierra norte del Ecuador. Ecuador: Universidad San Francisco de Quito, Colegio de Agricultura, Alimentos y Nutrición, Quito.
- Andersen RN, Walker HL. 1985. *Colletotrichum coccodes*: A pathogen of eastern black nightshade (*Solanum ptycanthum*). Weed Sci. 33: 902-905.
- Angulo R, editor. 2005. Uchuva: el cultivo. Colombia: Universidad de Bogotá Jorge Tadeo Lozano. 78 p.
- Arnold AE, Lutzoni F. 2007. Diversity and host range of foliar endophytes: are tropical leaves biodiversity hotspots? Ecology. 88: 541-549.
- Bruez E, Vallance J, Gerbore J, Lecomte P, Da Costa J, Guerin-Dubrana L, Rey P. 2014. Analyses of the temporal dynamics of fungal communities colonizing the healthy wood tissues of escaleaf-symptomatic and asymptomatic vines. PLoS One. 9: e95928.
- Catal M. 2002. Development and testing of oligonucleotide probes for detection and identification of some fungal pathogens and endophytes of conifers. PhD dissertation. USA: Michigan State University, Department of Botany and Plant Pathology.
- Cornejo-Espinoza FJ. 2014. Diversidad de hongos endófilos relacionados con lesiones necróticas en fréjol común (*Phaseolus vulgaris* L.) en la provincia de Loja, Ecuador. Bachelor's thesis. Ecuador: Pontificia Universidad Católica del Ecuador.
- Crous PW, Petrini O, Marais GF, Pretorius ZA, Rehder F. 1995. Occurrence of fungal endophytes in cultivars of *Triticum aestivum* in South Africa. Mycoscience. 36: 105-111.
- Crous PW, Verkley GJ, Groenewald E, Samson RA. 2009. CBS laboratory manual series 1: fungal biodiversity. The Netherlands, Central Bureau voor Schimmelcultures.
- Crozier J, Thomas SE, Aime MC, Evans HC, Holmes KA. 2006. Molecular characterization of fungal endophytic morphospecies isolated from stems and pods of *Theobroma cacao*. Plant Pathol. 55: 783-791.
- Dávila MF, Chaves-Acuña W, Yáñez J. 2018. Fungal communities associated with symptomatic leaves of naranjilla (*Solanum quitoense* Lam.) in the Ecuadorian Amazon region. Rev Ecuat Med Cienc Biol. 39: 39-49.
- De la Rotta, M.C. 2014. Enfermedades de la uchuva (*Physalis peruviana* L.). In: Carvalho CP, Moreno DA, editors. *Physalis peruviana: fruta andina para el mundo*. Spain: Programa Iberoamericano de Ciencia y Tecnología para el Desarrollo – CYTED. p. 49-78.
- Douanla-Meli C, Langer E, Mouafo FT. 2013. Fungal endophyte diversity and community

patterns in healthy and yellowing leaves of Citrus limon. Fungal Ecol. 6: 212-222.

Evans HC, Reeder RH. 2000. Fungi associated with Eichhornia crassipes (water hyacinth) in the upper Amazon Basin and prospects for their use in biological control. In: Julien M, Hill MP, Center T, Ding J, editors. Proceedings of the Meeting of the Global Working Group for the Biological Control and Integrated Control of Water Hyacinth; Australia: Australian Centre for International Agricultural Research.

Fischer G. 2000. Ecophysiological aspects of fruit growing in tropical highlands. Acta Hort. 531: 91-98.

Fischer G, Miranda D. 2012. Uchuva (Physalis peruviana L.). In: Fischer G, editor. Manual para el cultivo de frutales en el trópico Colombia: Produmedios p. 851-873.

Fischer G, Almanza-Merchán P, Miranda D. 2014. Importancia y cultivo de la uchuva (Physalis peruviana L.). Rev Bras Frutic. 36: 1-15.

Gazis R, Chaverri P. 2010. Diversity of fungal endophytes in leaves and stems of wild rubber trees (Hevea brasiliensis) in Peru. Fungal Ecol. 3: 240-254.

Ghimire SR, Charlton ND, Bell JD, Krishnamurthy YL, Craven KD. 2011. Biodiversity of fungal endophyte communities inhabiting switchgrass (Panicum virgatum L.) growing in the native tallgrass prairie of northern Oklahoma. Fungal Divers. 47: 19-27.

Gonzaga LL, Costa LEO, Santos TT, Araújo EF, Queiroz MV. 2015. Endophytic fungi from the genus Colletotrichum are abundant in the Phaseolus vulgaris and have high genetic diversity. J Appl Microbiol. 118: 485-496.

Hammer Ø, Harper DAT, Ryan PD. 2001. PAST: Paleontological statistics software package for education and data analysis. Palaeontol Electron. 4: 9.

Herrmann LW, Poitevin CG, Schuindt LC, De Almeida AB, Pimentel IC. 2019. Diversity of fungal endophytes in leaves of Mentha piperita and Mentha canadensis. Int J Botany Stud. 4: 44-49.

Hsieh HM, Lin CR, Fang MJ, Rogers JD, Fournier J, Lechat C, Ju YM. 2010. Phylogenetic status of Xylaria subgenus Pseudoxylaria among taxa

of the subfamily Xylarioideae (Xylariaceae) and phylogeny of the taxa involved in the subfamily. Mol Phylogenetics Evol. 54: 957-969.

Hyde KD, Soyong K. 2008. The fungal endophyte dilemma. Fungal Divers. 33: 163-173.

Kaur A, Oberhofer M, Juzumaitė M, Raja HA, Gulledge TV, Kao D, Faeth SH, Laster SM, Oberlies NH, Cech NB. 2016. Secondary metabolites from fungal endophytes of Echinacea purpurea suppress cytokine secretion by macrophage-type cells. Nat Prod Commun. 11: 1143.

Kernaghan G, Mayerhofer M, Griffin A. 2017. Fungal endophytes of wild and hybrid Vitis leaves and their potential for vineyard biocontrol. Can J Microbiol. 63: 583-95.

Kriel WM, Swart WJ, Crous PW. 2000. Foliar endophytes and their interactions with host plants, with specific reference to the Gymnospermae. In: Callow JA, editor. Advances in botanical research. London: Academic Press. p. 1-34.

Legge A. 1974. Notes on the history, cultivation and uses of Physalis peruviana L. J Roy Hort Soc. 99: 310-314.

Ling KS, Wechter WP, Somai BM, Walcott RR, Keinath AP. 2010. An improved real-time PCR system for broad-spectrum detection of Didymella bryoniae, the causal agent of gummy stem blight of cucurbits. Seed Sci Technol. 38: 692-703.

Loro M, Valero-Jiménez CA, Nozawa S, Márquez LM. 2012. Diversity and composition of fungal endophytes in semiarid Northwest Venezuela. J Arid Environ. 85: 46-55.

Maher N, Piot J, Bastien S, Vallance J, Rey P, Guérin-Dubrana L. 2012. Wood necrosis in esca-affected vines: types, relationships and possible links with foliar symptom expression. J Int Sci Vigne Vin. 46: 15-27.

Manamgoda DS, Rossman AY, Castlebury LA, Crous PW, Madrid H, Chukeatiro E, Hyde KD. 2014. The genus bipolaris. Stud Mycol. 79: 221-88.

Mazorra M. 2006. Aspectos anatómicos de la formación y crecimiento del fruto de uchuva Physalis peruviana (Solanaceae). Acta Biolo Colomb. 11: 69-81.

- McCain R. 1993. Goldenberry, passionfruit and white sapote: potential fruits for cool subtropical areas. In Janick J, Simon JE, editors. *New Crops*. New York: Wiley and Sons. p. 479-486.
- Miles LA, Lopera CA, González S, de García MC, Franco AE, Restrepo S. 2012. Exploring the biocontrol potential of fungal endophytes from an Andean Colombian Paramo ecosystem. *BioControl*. 57: 697-710.
- Mondal SN, Vicent A, Reis RF, Timmer LW. 2007. Saprophytic colonization of citrus twigs by *Diaporthe citri* and factors affecting pycnidial production and conidial survival. *Plant Dis*. 91: 387-392.
- Morton JF. 1987. Cape gooseberry. In: Morton JF, editor. *Fruits of warm climates*. Winterville, NC: Creative Resource Systems Inc. p. 430-434.
- Moya-Maldonado, A.E. 2016. Biodiversidad fúngica endófitas y epífitas de *Citrus sinensis*, naranjo dulce, de dos localidades de la Región Litoral del Ecuador. Bachelor's thesis. Ecuador: Pontificia Universidad Católica del Ecuador.
- Muñoz L. 2003. Análisis económico y financiero de la comercialización y producción de la uvilla como alternativa para la exportación. Bachelor's thesis. Ecuador: Escuela Superior Politécnica del Litoral, Guayaquil.
- Nilsson RH, Kristiansson E, Ryberg M, Hallenberg N, Larsson KH. 2008. Intraspecific ITS variability in the kingdom Fungi as expressed in the international sequence databases and its implications for molecular species identification. *Evol Bioinform*. 4: 193-201.
- Novoa R, Bojacá M, Galvis J, Fischer G. 2006. La madurez del fruto y el secado del cáliz influyen en el comportamiento poscosecha de la uchuva, almacenada a 12 °C (*Physalis peruviana* L.). *Agron Colomb*. 24: 77-86.
- Pacin AM, Gonzalez HHL, Etcheverry M, Resnik SL, Vivas L, Espin S. 2003. Fungi associated with food and feed commodities from Ecuador. *Mycopathologia*. 156: 87-92.
- Panno L, Bruno M, Voyron S, Anastasi A, Gnani G, Miserere L, Varese GC. 2013. Diversity, ecological role and potential biotechnological applications of marine fungi associated to the seagrass *Posidonia oceanica*. *N Biotechnol*. 30: 685-94.
- Paparu P, Dubois T, Gold CS, Niere B, Adipala E, Coyne D. 2008. Screenhouse and field persistence of nonpathogenic endophytic *Fusarium oxysporum* in *Musa* tissue culture plants. *Microb Ecol*. 55: 561-568.
- Parsa S, García-Lemos AM, Castillo K, Ortiz V, López-Lavalle LAB, Braun J, Vega FE. 2016. Fungal endophytes in germinated seeds of the common bean, *Phaseolus vulgaris*. *Fungal Biol*. 120: 783-790.
- Petrini O. 1996. Ecological and physiological aspects of host specificity in endophytic fungi. In: Redlin SC, Carris LM, editors. *Endophytic fungi of grasses and woody plants*. St Paul, MN: American Phytopathological Society. p. 87-100.
- Photita W, Lumyong S, Lumyong P, McKenzie EHC, Hyde KD. 2004. Are some endophytes of *Musa acuminata* latent pathogens? *Fungal Divers*. 16: 131-140.
- Prihastuti H, Cai L, Chen H, Hyde KD. 2009. Characterization of *Colletotrichum* species associated with coffee berries in Chiang Mai, Thailand. *Fungal Divers*. 39: 89-109.
- Promptutha L, Lumyong S, Dhanasekaran V, McKenzie E, Hyde K, Jeewon R. 2007. A phylogenetic evaluation of whether endophytes become saprotrophs at host senescence. *Microb Ecol*. 53: 579-590.
- Puente LA, Pinto-Muñoz CA, Castro ES, Cortés M. 2011. *Physalis peruviana* Linnaeus, the multiple properties of a highly functional fruit: A review. *Food Res Int*. 44: 1733-1740.
- Raja HA, Kaur A, El-Elmag T, Figueroa M, Kumar R, Deep G, Agarwal R, Faeth SH, Cech NB, Oberlies NH. 2015. Phylogenetic and chemical diversity of fungal endophytes isolated from *Silybum marianum* (L.) Gaertn. (milk thistle). *Mycology*. 6: 8-27.
- Ramadan MF, Mörsel JT. 2003. Oil goldenberry (*Physalis peruviana* L.). *J Agric Food Chem*. 51: 969-974.
- Ramírez J, Fernandez ED, Rodolfi M, Solveig T. 2006. Actividad antagónica de hongos endófitos de plantas medicinales del Ecuador sobre bacterias patógenas. *Bol Micol*. 21: 49-53.

- Rao VG, Subramoniam V. 1976. A new post-harvest disease of cape-gooseberry. *J Univ Bombay*. 72:58-61.
- Rehm S, Espig G. 1991. Fruit. In: Sigmund R, Gustav E, editors. *The cultivated plants of the tropics and subtropics, cultivation, economic value, utilization*. Germany: Verlag Josef Margraf. p. 169-245.
- Rekab D, Del Sorbo G, Reggio C, Zoina A, Firrao G. 2004. Polymorphisms in nuclear rDNA and mtDNA reveal the polyphyletic nature of isolates of *Phomopsis* pathogenic to sunflower and a tight monophyletic clade of defined geographic origin. *Mycol Res*. 108:393-402.
- Rocha DOL, Reis GM, da Silva VN, Braghini R, Teixeira MM, Correa B. 2011. Molecular characterization and fumonisin production by *Fusarium verticillioides* isolated from corn grains of different geographic origins in Brazil. *Int J Food Microbiol*. 145: 9-21.
- Rodrigues A, Mueller UG, Ishak HD, Bacci Jr M, Pagnocca FC. 2011. Ecology of microfungal communities in gardens of fungus-growing ants (Hymenoptera: Formicidae): a year-long survey of three species of attine ants in Central Texas. *FEMS Microbiol Ecol*. 78: 244-55.
- Rodríguez R, Redman R. 2008. More than 400 million years of evolution and some plants still can't make it on their own: plant stress tolerance via fungal symbiosis. *J Exp Bot*. 59: 1109-1114.
- Rodríguez-Amézquita JE, Velandia-Monsalve J, Viteri-Rosero SE. 2010. Evaluación de microorganismos aislados de gallinaza por su potencial para el biocontrol de fusarium (*F. oxysporum*) en plántulas de uchuva (*Physalis peruviana*). *Rev Fac Nal Agr Medellín*. 63: 5499-5509.
- Schulz B, Boyle C. 2006. What are endophytes? In: Schulz B, Boyle CJ, Sieber TN, editors. *Microbial root endophytes*. Berlin: Springer. p. 1-13.
- Shrestha P, Szaro TM, Bruns TD, Taylor JW. 2011. Systematic search for cultivatable fungi that best deconstruct cell walls of *Miscanthus* and sugarcane in the field. *Appl Environ Microbiol*. 77: 5490-5504.
- Sieber T. 2007. Endophytic fungi in forest trees: are they mutualists? *Fungal Biol Rev*. 21: 75-89.
- Stefańczyk E, Sobkowiak S, Brylińska M, Śliwka J. 2016. Diversity of *Fusarium* spp. associated with dry rot of potato tubers in Poland. *Eur J Plant Pathol*. 145: 871-884.
- Sun J, Guo L, Zang W, Ping W, Chi D. 2008. Diversity and ecological distribution of endophytic fungi associated with medicinal plants. *Sci China C Life Sci*. 51: 751-759.
- Thomas SE, Crozier J, Aime MC, Evans HC, Holmes KA. 2008. Molecular characterization of fungal endophytic morphospecies associated with the indigenous forest tree, *Theobroma gileri*, in Ecuador. *Mycol Res*. 112: 852-860.
- Thomas DC, Vandegrift R, Ludden A, Carroll GC, Roy BA. 2016. Spatial ecology of the fungal genus *Xylaria* in a tropical cloud forest. *Biotropica*. 48: 381-393.
- Van Bael SA, Fernández-Marín H, Valencia MC, Rojas EI, Wcislo WT, Herre EA. 2009. Two fungal symbioses collide: endophytic fungi are not welcome in leaf-cutting ant gardens. *Proc R Soc Lond B Biol Sci*. 276:2419-2426.
- Videira SI, Groenewald JZ, Braun U, Shin HD, Crous PW. 2016. All that glitters is not *Ramularia*. *Stud Mycol*. 83: 49-163.
- Waculicz-Andrade CE, Savi DC, Bini AP, Adamoski D, Goulin EH, Silva GJ, Massola NS, Terasawa LG, Kava V, Glienke C. 2017. *Colletotrichum gloeosporioides sensu stricto*: an endophytic species or citrus pathogen in Brazil?. *Australas Plant Pathol*. 46: 191-203.
- Weir BS, Johnston PR, Damm U. 2012. The *Colletotrichum gloeosporioides* species complex. *Stud Mycology*. 73: 115-80.
- White TF, Brunus T, Lee S, Taylor J. 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky FS, White TT, editors. *PCR protocol: a guide to methods and applications*. San Diego: Academic Press. p. 315-322.
- Zapata JL, Saldarriaga A, Londoño M, Díaz C. 2002. Manejo del cultivo de la uchuva en Colombia. Colombia: Corpoica. Boletín técnico.

